arXiv:1703.07861v1 [math.AP] 22 Mar 2017

A FRACTIONAL KIRCHHOFF PROBLEM INVOLVING A SINGULAR TERM AND A CRITICAL NONLINEARITY

ALESSIO FISCELLA

Abstract. In this paper we consider the following critical nonlocal problem

$$\begin{cases} M\left(\iint\limits_{\mathbb{R}^{2N}} \frac{|u(x)-u(y)|^2}{|x-y|^{N+2s}}dxdy\right)(-\Delta)^s u = \frac{\lambda}{u^\gamma} + u^{2_s^*-1} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases}$$

where Ω is an open bounded subset of $\mathbb{R}^N$ with continuous boundary, dimension $N > 2s$ with parameter $s \in (0,1)$, $2_s^* = 2N/(N-2s)$ is the fractional critical Sobolev exponent, $\lambda > 0$ is a real parameter, exponent $\gamma \in (0,1)$, M models a Kirchhoff type coefficient, while $(-\Delta)^s$ is the fractional Laplace operator. In particular, we cover the delicate degenerate case, that is when the Kirchhoff function M is zero at zero. By combining variational methods with an appropriate truncation argument, we provide the existence of two solutions.

1. Introduction

This paper is devoted to the study of a class of Kirchhoff type problems driven by a nonlocal fractional operator and involving a singular term and a critical nonlinearity. More precisely, we consider

$$\begin{cases} \left(\iint\limits_{\mathbb{R}^{2N}} \frac{|u(x)-u(y)|^2}{|x-y|^{N+2s}}dxdy\right)^{\theta-1}(-\Delta)^s u = \frac{\lambda}{u^\gamma} + u^{2_s^*-1} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases} \tag{1.1}$$

where Ω is an open bounded subset of $\mathbb{R}^N$ with continuous boundary, dimension $N > 2s$ with parameter $s \in (0,1)$, $2_s^* = 2N/(N-2s)$ is the fractional critical Sobolev exponent, $\lambda > 0$ is a real parameter, exponent $\theta \in (1, 2_s^*/2)$ while $\gamma \in (0,1)$. Here $(-\Delta)^s$ is the fractional Laplace operator defined, up to normalization factors, by the Riesz potential as

$$(-\Delta)^s \varphi(x) = \int_{\mathbb{R}^N} \frac{2\varphi(x) - \varphi(x+y) - \varphi(x-y)}{|y|^{N+2s}}dy, \quad x \in \mathbb{R}^N,$$

along any $\varphi \in C_0^\infty(\Omega)$; we refer to [11] and the recent monograph [22] for further details on the fractional Laplacian and the fractional Sobolev spaces $H^s(\mathbb{R}^N)$ and $H_0^s(\Omega)$.

Key words and phrases. Kirchhoff type problems, fractional Laplacian, singularities, critical nonlinearities, perturbation methods.

2010 AMS Subject Classification. Primary: 35J75, 35R11, 49J35; Secondary: 35A15, 45G05, 35S15.

As well explained in [11], problem (1.1) is the fractional version of the following nonlinear problem

$$\begin{cases} -M\left(\int_\Omega |\nabla u(x)|^2 dx\right)\Delta u = \dfrac{\lambda}{u^\gamma} + u^{2^*-1} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{in } \partial\Omega, \end{cases} \tag{1.2}$$

where Δ denotes the classical Laplace operator while, just for a general discussion, $M(t) = t^{\theta-1}$ for any $t \in \mathbb{R}_0^+$. In literature, problems like (1.1) and (1.2) are called of Kirchhoff type whenever the function $M : \mathbb{R}_0^+ \to \mathbb{R}_0^+$ models the Kirchhoff prototype, given by

$$M(t) = a + bt^{\theta-1}, \quad a, b \geq 0, \ a + b > 0, \ \theta \geq 1. \tag{1.3}$$

In particular, when $M(t) \geq$ *constant* > 0 for any $t \in \mathbb{R}_0^+$, Kirchhoff problems are said to be *non–degenerate* and this happens for example if $a > 0$ in the model case (1.3). While, if $M(0) = 0$ but $M(t) > 0$ for any $t \in \mathbb{R}^+$, Kirchhoff problems are called *degenerate*. Of course, for (1.3) this occurs when $a = 0$.

This kind of nonlocal problems has been widely studied in recent years. We refer to [17, 18, 19, 20, 21] for different Kirchhoff problems with M like in (1.3), driven by the Laplace operator and involving a singular term of type $u^{-\gamma}$. In [20], the authors study a Kirchhoff problem with a singular term and a Hardy potential, by using the Nehari method. The same approach is used in [19] for a singular Kirchhoff problem with also a subcritical term. In [17], strongly assuming $a > 0$ in (1.3), the authors prove the existence of two solutions for a Kirchhoff problem like (1.2), by combining perturbation and variational methods. While in [18], they provide a uniqueness result for a singular Kirchhoff problem involving a negative critical nonlinearity, by a minimization argument. By arguing similarly to [17], the authors of [21] give the existence of two solutions for a critical Kirchhoff problem with a singular term of type $|x|^{-\beta}u^{-\gamma}$.

Problem (1.1) has been studied in [3] when $\theta = 1$, namely without a Kirchhoff coefficient. In [3], they prove the existence of two solutions by applying the sub/supersolutions and Sattinger methods. In [8], the authors generalize the results of Section 3 of [3] to the delicate case of the p–fractional Laplace operator $(-\Delta)_p^s$. While in the last section of [1], the authors provide the existence of a solution for nonlinear fractional problems with a singularity like $u^{-\gamma}$ and a fractional Hardy term, by perturbation methods. Concerning fractional Kirchhoff problems involving critical nonlinearities, we refer to [2, 9, 13, 14, 16, 23] for existence results and to [5, 12, 24, 25, 29] for multiplicity results. In particular, in [9, 13, 14, 23] different singular terms appear, but given by the fractional Hardy potential.

Inspired by the above works, we study a multiplicity result for problem (1.1). As far as we know, a fractional Kirchhoff problem involving a singular term of type $u^{-\gamma}$ has not been studied yet. Thus, we can state our result as follows.

Theorem 1.1. *Let $s \in (0,1)$, $N > 2s$, $\theta \in (1, 2_s^*/2)$, $\gamma \in (0,1)$ and let Ω be an open bounded subset of $\mathbb{R}^N$ with $\partial\Omega$ continuous. Then, there exists $\overline{\lambda} > 0$ such that for any $\lambda \in (0, \overline{\lambda})$ problem (1.1) has at least two different solutions.*

The first solution of problem (1.1) is obtained by a suitable minimization argument, where we must pay attention to the nonlocal nature of the fractional Laplacian. Concerning the second solution, because of the presence of $u^{-\gamma}$, we can not apply the usual critical point theory to problem (1.1). For this, we first study a perturbed problem obtained truncating the singular term $u^{-\gamma}$. Then, by approximation we get our second solution of (1.1).

Finally, we observe that Theorem 1.1 generalizes in several directions the first part of [3, Theorem 4.1] and [17, Theorem 1.1].

The paper is organized as follows. In Section 2, we discuss the variational formulation of problem (1.1) and we introduce the perturbed problem. In Section 3, we prove the existence of the first solution of (1.1) and we give a possible generalization of this existence result, at the end of the section. In Section 4, we prove the existence of a mountain pass solution for the perturbed problem. In Section 5, we prove Theorem 1.1.

2. Variational setting

Throughout the paper we assume that $s \in (0,1)$, $N > 2s$, $\theta \in (1, 2_s^*/2)$, $\gamma \in (0,1)$ *and* Ω *is an open bounded subset of* $\mathbb{R}^N$ *with* $\partial\Omega$ *continuous, without further mentioning. As a matter of notations, we denote with* $\varphi^+ = \max\{\varphi, 0\}$ *and* $\varphi^- = \max\{-\varphi, 0\}$ *respectively the positive and negative part of a function* φ.

Problem (1.1) has a variational structure and the natural space where finding solutions is the homogeneous fractional Sobolev space $H_0^s(\Omega)$. In order to study (1.1), it is important to encode the "boundary condition" $u = 0$ in $\mathbb{R}^N \setminus \Omega$ in the weak formulation, by considering also that the interaction between Ω and its complementary in $\mathbb{R}^N$ gives a positive contribution in the so called *Gagliardo norm*, given as

$$\|u\|_{H^s(\mathbb{R}^N)} = \|u\|_{L^2(\mathbb{R}^N)} + \Big(\iint\limits_{\mathbb{R}^{2N}} \frac{|u(x)-u(y)|^2}{|x-y|^{N+2s}} dxdy \Big)^{1/2}. \tag{2.1}$$

The functional space that takes into account this boundary condition will be denoted by X_0 and it is defined as

$$X_0 = \big\{ u \in H^s(\mathbb{R}^N) : \ u = 0 \text{ a.e. in } \mathbb{R}^N \setminus \Omega \big\}.$$

We refer to [26] for a general definition of X_0 and its properties. We also would like to point out that, when $\partial\Omega$ is continuous, by [15, Theorem 6] the space $C_0^\infty(\Omega)$ is dense in X_0, with respect to the norm (2.1). This last point will be used to overcome the singularity in problem (1.1).

In X_0 we can consider the following norm

$$\|u\|_{X_0} = \Big(\iint\limits_{\mathbb{R}^{2N}} \frac{|u(x)-u(y)|^2}{|x-y|^{N+2s}} dxdy \Big)^{1/2},$$

which is equivalent to the usual one defined in (2.1) (see [26, Lemma 6]). We also recall that $(X_0, \|\cdot\|_{X_0})$ is a Hilbert space, with the scalar product defined as

$$\langle u, v \rangle_{X_0} = \iint\limits_{\mathbb{R}^{2N}} \frac{(u(x)-u(y))(v(x)-v(y))}{|x-y|^{N+2s}} dxdy.$$

From now on, in order to simplify the notation, we will denote $\|\cdot\|_{X_0}$ *and* $\langle\cdot,\cdot\rangle_{X_0}$ *by* $\|\cdot\|$ *and* $\langle\cdot,\cdot\rangle$ *respectively, and* $\|\cdot\|_{L^q(\Omega)}$ *by* $\|\cdot\|_q$ *for any* $q \in [1,\infty]$.

In order to present the weak formulation of (1.1) and taking into account that we are looking for positive solutions, we will consider the following Kirchhoff problem

$$\begin{cases} \Big(\iint\limits_{\mathbb{R}^{2N}} \frac{|u(x)-u(y)|^2}{|x-y|^{N+2s}} dxdy \Big)^{\theta-1} (-\Delta)^s u = \dfrac{\lambda}{(u^+)^\gamma} + (u^+)^{2_s^*-1} & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega. \end{cases} \tag{2.2}$$

We say that $u \in X_0$ is a (weak) solution of problem (2.2), if u satisfies

$$\|u\|^{2(\theta-1)} \langle u, \varphi \rangle = \lambda \int_\Omega \frac{\varphi}{(u^+)^\gamma} dx + \int_\Omega (u^+)^{2_s^*-1} \varphi, \tag{2.3}$$

for any $\varphi \in X_0$. Problem (2.2) has a variational structure and $J_\lambda : X_0 \to \mathbb{R}$, defined by

$$J_\lambda(u) = \frac{1}{2\theta}\|u\|^{2\theta} - \frac{\lambda}{1-\gamma}\int_\Omega (u^+)^{1-\gamma}dx - \frac{1}{2_s^*}\|u^+\|_{2_s^*}^{2_s^*},$$

is the underlying functional associated to (2.2). Because of the presence of a singular term in (2.2), functional J_λ is not differentiable on X_0. Therefore, we can not apply directly the usual critical point theory to J_λ, in order to solve problem (2.2). However, it is possible to find a first solution of (2.2) by using a local minimization argument. In order to get the second solution of (2.2), we have to study an associated approximating problem. That is, for any $n \in \mathbb{N}$, we consider the following perturbed problem

$$\begin{cases} \left(\iint\limits_{\mathbb{R}^{2N}} \frac{|u(x)-u(y)|^2}{|x-y|^{N+2s}}dxdy\right)^{\theta-1} (-\Delta)^s u = \frac{\lambda}{(u^+ + 1/n)^\gamma} + (u^+)^{2_s^*-1} & \text{in } \Omega, \\ u = 0 \quad \text{in } \mathbb{R}^N \setminus \Omega. \end{cases} \tag{2.4}$$

For this, we say that $u \in X_0$ is a (weak) solution of problem (2.4), if u satisfies

$$\|u\|^{2(\theta-1)}\langle u, \varphi\rangle = \lambda\int_\Omega \frac{\varphi}{(u^+ + 1/n)^\gamma}dx + \int_\Omega (u^+)^{2_s^*-1}\varphi, \tag{2.5}$$

for any $\varphi \in X_0$. In this case, solutions of (2.4) correspond to the critical points of functional $J_{n,\lambda} : X_0 \to \mathbb{R}$, set as

$$J_{n,\lambda}(u) = \frac{1}{2\theta}\|u\|^{2\theta} - \frac{\lambda}{1-\gamma}\int_\Omega [(u^+ + 1/n)^{1-\gamma} - (1/n)^{1-\gamma}]dx - \frac{1}{2_s^*}\|u^+\|_{2_s^*}^{2_s^*}. \tag{2.6}$$

It is immediate to see that $J_{n,\lambda}$ is of class $C^1(X_0)$.

We conclude this section recalling the best constant of the fractional Sobolev embedding, which will be very useful to study the compactness property of functional $J_{n,\lambda}$. That is, we consider

$$S = \inf_{\substack{v\in H^s(\mathbb{R}^N)\\ v\neq 0}} \frac{\iint\limits_{\mathbb{R}^{2N}} \frac{|v(x)-v(y)|^2}{|x-y|^{N+2s}}dxdy}{\left(\int_{\mathbb{R}^N}|v(x)|^{2_s^*}dx\right)^{2/2_s^*}}, \tag{2.7}$$

which is well defined and strictly positive, as shown in [10, Theorem 1.1].

3. A first solution for problem (1.1)

In this section we prove the existence of a solution for problem (1.1) by a local minimization argument. For this, we first study the geometry of functional J_λ.

Lemma 3.1. *There exist numbers $\rho \in (0,1]$, $\lambda_0 = \lambda_0(\rho) > 0$ and $\alpha = \alpha(\rho) > 0$ such that $J_\lambda(u) \geq \alpha$ for any $u \in X_0$, with $\|u\| = \rho$, and for any $\lambda \in (0, \lambda_0]$.*

Furthermore, set

$$m_\lambda = \inf\left\{J_\lambda(u) :\ u \in \overline{B}_\rho\right\},$$

where $\overline{B}_\rho = \{u \in X_0 :\ \|u\| \leq \rho\}$. Then, $m_\lambda < 0$ for any $\lambda \in (0, \lambda_0]$.

Proof. Let $\lambda > 0$. From the Hölder inequality and (2.7), for any $u \in X_0$ we have

$$\int_\Omega (u^+)^{1-\gamma}dx \leq |\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}}\|u\|_{2_s^*}^{1-\gamma} \leq |\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}} S^{-\frac{1-\gamma}{2}}\|u\|^{1-\gamma}. \tag{3.1}$$

Hence, by using again (2.7) and (3.1) we get

$$J_\lambda(u) \geq \frac{1}{2\theta}\|u\|^{2\theta} - \frac{S^{-\frac{2_s^*}{2}}}{2_s^*}\|u\|^{2_s^*} - \frac{\lambda}{1-\gamma}|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}} S^{-\frac{1-\gamma}{2}}\|u\|^{1-\gamma}.$$

Since $1-\gamma < 1 < 2\theta < 2_s^*$, the function

$$\eta(t) = \frac{1}{2\theta}t^{2\theta-1+\gamma} - \frac{S^{-\frac{2_s^*}{2}}}{2_s^*}t^{2_s^*-1+\gamma}, \quad t \in [0,1]$$

admits a maximum at some $\rho \in (0,1]$ small enough, that is $\max_{t\in[0,1]} \eta(t) = \eta(\rho) > 0$. Thus, let

$$\lambda_0 = \frac{(1-\gamma)S^{\frac{1-\gamma}{2}}}{2|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}}}\eta(\rho),$$

then for any $u \in X_0$ with $\|u\| = \rho \leq 1$ and for any $\lambda \leq \lambda_0$, we get $J_\lambda(u) \geq \rho^{1-\gamma}\eta(\rho)/2 = \alpha > 0$.

Furthermore, fixed $v \in X_0$ with $v^+ \not\equiv 0$, for $t \in (0,1)$ sufficiently small

$$J_\lambda(tv) = \frac{t^{2\theta}}{2\theta}\|v\|^{2\theta} - t^{1-\gamma}\frac{\lambda}{1-\gamma}\int_\Omega (v^+)^{1-\gamma}dx - \frac{t^{2_s^*}}{2_s^*}\|v^+\|_{2_s^*}^{2_s^*} < 0,$$

being $1-\gamma < 1 < 2\theta < 2_s^*$. This concludes the proof. □

We are now ready to prove the existence of the first solution of (1.1).

Theorem 3.2. *Let λ_0 be given as in Lemma 3.1. Then, for any $\lambda \in (0, \lambda_0]$ problem (1.1) has a solution $u_0 \in X_0$, with $J_\lambda(u_0) < 0$.*

Proof. Fix $\lambda \in (0, \lambda_0]$ and let ρ be as given in Lemma 3.1. We first prove there exists $u_0 \in \overline{B}_\rho$ such that $J_\lambda(u_0) = m_\lambda < 0$. Let $\{u_k\}_k \subset \overline{B}_\rho$ be a minimizing sequence for m_λ, that is such that

$$\lim_{k\to\infty} J_\lambda(u_k) = m_\lambda. \tag{3.2}$$

Since $\{u_k\}_k$ is bounded in X_0, by applying [26, Lemma 8] and [6, Theorem 4.9], there exist a subsequence, still denoted by $\{u_k\}_k$, and a function $u_0 \in \overline{B}_\rho$ such that, as $k \to \infty$ we have

$$\begin{aligned} &u_k \rightharpoonup u_0 \text{ in } X_0, && u_k \rightharpoonup u_0 \text{ in } L^{2_s^*}(\Omega),\\ &u_k \to u_0 \text{ in } L^p(\Omega) \text{ for any } p \in [1, 2_s^*), && u_k \to u_0 \text{ a.e. in } \Omega. \end{aligned} \tag{3.3}$$

Since $\gamma \in (0,1)$, by the Hölder inequality, for any $k \in \mathbb{N}$ we have

$$\left|\int_\Omega (u_k^+)^{1-\gamma}dx - \int_\Omega (u_0^+)^{1-\gamma}dx\right| \leq \int_\Omega \left|u_k^+ - u_0^+\right|^{1-\gamma}dx \leq \|u_k^+ - u_0^+\|_2^{1-\gamma}|\Omega|^{\frac{1+\gamma}{2}},$$

which yields, by (3.3)

$$\lim_{k\to\infty}\int_\Omega (u_k^+)^{1-\gamma}dx = \int_\Omega (u_0^+)^{1-\gamma}dx. \tag{3.4}$$

Let $w_k = u_k - u_0$, by [7, Theorem 2] it holds true that

$$\|u_k\|^2 = \|w_k\|^2 + \|u_0\|^2 + o(1), \quad \|u_k\|_{2_s^*}^{2_s^*} = \|w_k\|_{2_s^*}^{2_s^*} + \|u_0\|_{2_s^*}^{2_s^*} + o(1) \tag{3.5}$$

as $k \to \infty$. Since $\{u_k\}_k \subset \overline{B}_\rho$, by (3.5) for k sufficiently large, we have $w_k \in \overline{B}_\rho$. Lemma 3.1 implies that for any $u \in X_0$, with $\|u\| = \rho$, we get

$$\frac{1}{2\theta}\|u\|^{2\theta} - \frac{1}{2_s^*}\|u^+\|_{2_s^*}^{2_s^*} \geq \alpha > 0,$$

and from this, being $\rho \leq 1$, for k sufficiently large we have

$$\frac{1}{2\theta}\|w_k\|^{2\theta} - \frac{1}{2_s^*}\|w_k^+\|_{2_s^*}^{2_s^*} > 0. \tag{3.6}$$

Thus, by (3.2), (3.4)–(3.6) and considering $\theta \geq 1$, it follows that as $k \to \infty$

$$\begin{aligned} m_\lambda &= J_\lambda(u_k) + o(1) \\ &= \frac{1}{2\theta}\left(\|w_k\|^2 + \|u_0\|^2\right)^\theta - \frac{\lambda}{1-\gamma}\int_\Omega (u_0^+)^{1-\gamma}dx - \frac{1}{2_s^*}\left(\|w_k^+\|_{2_s^*}^{2_s^*} + \|u_0^+\|_{2_s^*}^{2_s^*}\right) + o(1) \\ &\geq J_\lambda(u_0) + \frac{1}{2\theta}\|w_k\|^{2\theta} - \frac{1}{2_s^*}\|w_k^+\|_{2_s^*}^{2_s^*} + o(1) \geq J_\lambda(u_0) + o(1) \geq m_\lambda, \end{aligned}$$

being $u_0 \in \overline{B}_\rho$. Hence, u_0 is a local minimizer for J_λ, with $J_\lambda(u_0) = m_\lambda < 0$ which implies that u_0 is nontrivial.

Now, we prove that u_0 is a positive solution of (2.2). For any $\psi \in X_0$, with $\psi \geq 0$ a.e. in $\mathbb{R}^N$, let us consider a $t > 0$ sufficiently small so that $u_0 + t\psi \in \overline{B}_\rho$. Since u_0 is a local minimizer for J_λ, we have

$$\begin{aligned} 0 &\leq J_\lambda(u_0 + t\psi) - J_\lambda(u_0) \\ &= \frac{1}{2\theta}\left(\|u_0 + t\psi\|^{2\theta} - \|u_0\|^{2\theta}\right) - \frac{\lambda}{1-\gamma}\int_\Omega \left[((u_0 + t\psi)^+)^{1-\gamma} - (u_0^+)^{1-\gamma}\right]dx \\ &\quad - \frac{1}{2_s^*}\left(\|u_0 + t\psi\|_{2_s^*}^{2_s^*} - \|u_0^+\|_{2_s^*}^{2_s^*}\right). \end{aligned}$$

From this, dividing by $t > 0$ and passing to the limit as $t \to 0^+$, it follows that

$$\liminf_{t\to 0^+} \frac{\lambda}{1-\gamma}\int_\Omega \frac{((u_0 + t\psi)^+)^{1-\gamma} - (u_0^+)^{1-\gamma}}{t}dx \leq \|u_0\|^{2(\theta-1)}\langle u_0, \psi\rangle - \int_\Omega (u_0^+)^{2_s^*-1}\psi dx. \tag{3.7}$$

We observe that

$$\frac{1}{1-\gamma}\cdot\frac{((u_0 + t\psi)^+)^{1-\gamma} - (u_0^+)^{1-\gamma}}{t} = ((u_0 + \xi t\psi)^+)^{-\gamma}\psi \quad \text{a.e. in } \Omega,$$

with $\xi \in (0,1)$ and $((u_0 + \xi t\psi)^+)^{-\gamma} \to (u_0^+)^{-\gamma}\psi$ a.e. in Ω, as $t \to 0^+$. Thus, by the Fatou lemma, we obtain

$$\lambda\int_\Omega (u_0^+)^{-\gamma}\psi dx \leq \liminf_{t\to 0^+} \frac{\lambda}{1-\gamma}\int_\Omega \frac{((u_0 + t\psi)^+)^{1-\gamma} - (u_0^+)^{1-\gamma}}{t}dx. \tag{3.8}$$

Therefore, combining (3.7) and (3.8) we get

$$\|u_0\|^{2(\theta-1)}\langle u_0, \psi\rangle - \lambda\int_\Omega (u_0^+)^{-\gamma}\psi dx - \int_\Omega (u_0^+)^{2_s^*-1}\psi dx \geq 0, \tag{3.9}$$

for any $\psi \in X_0$, with $\psi \geq 0$ a.e. in $\mathbb{R}^N$.

Since $J_\lambda(u_0) < 0$ and by Lemma 3.1, we have $u_0 \in B_\rho$. Hence, there exists $\delta \in (0,1)$ such that $(1+t)u_0 \in \overline{B}_\rho$ for any $t \in [-\delta, \delta]$. Let us define $I_\lambda(t) = J_\lambda((1+t)u_0)$. Since u_0 is a local minimizer for J_λ in $\overline{B}_\rho$, functional I_λ has a minimum at $t = 0$, that is

$$I_\lambda'(0) = \|u_0\|^{2\theta} - \lambda\int_\Omega (u_0^+)^{1-\gamma}dx - \int_\Omega (u_0^+)^{2_s^*}dx = 0. \tag{3.10}$$

For any $\varphi \in X_0$ and any $\varepsilon > 0$, let us define $\psi_\varepsilon = u_0^+ + \varepsilon\varphi$. Then, by (3.9) we have

$$\begin{aligned} 0 &\le \|u_0\|^{2(\theta-1)}\langle u_0, \psi_\varepsilon^+\rangle - \lambda\int_\Omega (u_0^+)^{-\gamma}\psi_\varepsilon^+ dx - \int_\Omega (u_0^+)^{2_s^*-1}\psi_\varepsilon^+ dx \\ &= \|u_0\|^{2(\theta-1)}\langle u_0, \psi_\varepsilon + \psi_\varepsilon^-\rangle - \lambda\int_\Omega (u_0^+)^{-\gamma}(\psi_\varepsilon + \psi_\varepsilon^-)dx - \int_\Omega (u_0^+)^{2_s^*-1}(\psi_\varepsilon + \psi_\varepsilon^-)dx. \end{aligned} \tag{3.11}$$

We observe that, for a.e. x, $y \in \mathbb{R}^N$, we obtain

$$\begin{aligned} &(u_0(x) - u_0(y))(u_0^-(x) - u_0^-(y)) \\ &\quad = -u_0^+(x)u_0^-(y) - u_0^-(x)u_0^+(y) - \left[u_0^-(x) - u_0^-(y)\right]^2 \\ &\quad \le -\left|u_0^-(x) - u_0^-(y)\right|^2, \end{aligned} \tag{3.12}$$

from which we immediately get

$$(u_0(x) - u_0(y))(u_0^+(x) - u_0^+(y)) \le |u_0(x) - u_0(y)|^2.$$

From the last inequality, it follows that

$$\begin{aligned} \langle u_0, \psi_\varepsilon + \psi_\varepsilon^-\rangle &= \iint_{\mathbb{R}^{2N}} \frac{(u_0(x) - u_0(y))(\psi_\varepsilon(x) + \psi_\varepsilon^-(x) - \psi_\varepsilon(y) - \psi_\varepsilon^-(y))}{|x-y|^{N+2s}}dxdy \\ &\le \iint_{\mathbb{R}^{2N}} \frac{|u_0(x) - u_0(y)|^2}{|x-y|^{N+2s}}dxdy + \varepsilon\iint_{\mathbb{R}^{2N}} \frac{(u_0(x) - u_0(y))(\varphi(x) - \varphi(y))}{|x-y|^{N+2s}}dxdy \\ &\quad + \iint_{\mathbb{R}^{2N}} \frac{(u_0(x) - u_0(y))(\psi_\varepsilon^-(x) - \psi_\varepsilon^-(y))}{|x-y|^{N+2s}}dxdy. \end{aligned} \tag{3.13}$$

Hence, denoting with $\Omega_\varepsilon = \left\{x \in \mathbb{R}^N : \ u_0^+(x) + \varepsilon\varphi(x) \le 0\right\}$ and by combining (3.11) with (3.13), we get

$$\begin{aligned} 0 &\le \|u_0\|^{2\theta} + \varepsilon\|u_0\|^{2(\theta-1)}\langle u_0, \varphi\rangle + \|u_0\|^{2(\theta-1)}\langle u_0, \psi_\varepsilon^-\rangle \\ &\quad - \lambda\int_\Omega (u_0^+)^{-\gamma}(u_0^+ + \varepsilon\varphi)dx - \int_\Omega (u_0^+)^{2_s^*-1}(u_0^+ + \varepsilon\varphi)dx \\ &\quad + \lambda\int_{\Omega_\varepsilon} (u_0^+)^{-\gamma}(u_0^+ + \varepsilon\varphi)dx + \int_{\Omega_\varepsilon} (u_0^+)^{2_s^*-1}(u_0^+ + \varepsilon\varphi)dx \\ &\le \|u_0\|^{2\theta} - \lambda\int_\Omega (u_0^+)^{1-\gamma}dx - \int_\Omega (u_0^+)^{2_s^*}dx + \|u_0\|^{2(\theta-1)}\langle u_0, \psi_\varepsilon^-\rangle \\ &\quad + \varepsilon\left[\|u_0\|^{2(\theta-1)}\langle u_0, \varphi\rangle - \lambda\int_\Omega (u_0^+)^{-\gamma}\varphi dx - \int_\Omega (u_0^+)^{2_s^*-1}\varphi dx\right] \\ &= \|u_0\|^{2(\theta-1)}\langle u_0, \psi_\varepsilon^-\rangle + \varepsilon\left[\|u_0\|^{2(\theta-1)}\langle u_0, \varphi\rangle - \lambda\int_\Omega (u_0^+)^{-\gamma}\varphi dx - \int_\Omega (u_0^+)^{2_s^*-1}\varphi dx\right], \end{aligned} \tag{3.14}$$

where last equality follows from (3.10). Arguing similarly to (3.12), for a.e. x, $y \in \mathbb{R}^N$ we have

$$(u_0(x) - u_0(y))(u_0^+(x) - u_0^+(y)) \ge \left|u_0^+(x) - u_0^+(y)\right|^2. \tag{3.15}$$

Thus, denoting with

$$\mathcal{U}_\varepsilon(x,y) = \frac{(u_0(x) - u_0(y))(\psi_\varepsilon^-(x) - \psi_\varepsilon^-(y))}{|x-y|^{N+2s}},$$

by the symmetry of the fractional kernel and (3.15), we get

$$\begin{aligned}\langle u_0, \psi_\varepsilon^- \rangle &= \iint\limits_{\Omega_\varepsilon \times \Omega_\varepsilon} \mathcal{U}_\varepsilon(x,y)dxdy + 2 \iint\limits_{\Omega_\varepsilon \times (\mathbb{R}^N \setminus \Omega_\varepsilon)} \mathcal{U}_\varepsilon(x,y)dxdy \\ &\leq -\varepsilon \left(\iint\limits_{\Omega_\varepsilon \times \Omega_\varepsilon} \mathcal{U}(x,y)dxdy + 2 \iint\limits_{\Omega_\varepsilon \times (\mathbb{R}^N \setminus \Omega_\varepsilon)} \mathcal{U}(x,y)dxdy \right) \\ &\leq 2\varepsilon \iint\limits_{\Omega_\varepsilon \times \mathbb{R}^N} |\mathcal{U}(x,y)|\, dxdy,\end{aligned} \tag{3.16}$$

where we set

$$\mathcal{U}(x,y) = \frac{(u_0(x) - u_0(y))(\varphi(x) - \varphi(y))}{|x-y|^{N+2s}}.$$

Clearly $\mathcal{U} \in L^1(\mathbb{R}^{2N})$, so that for any $\sigma > 0$ there exists R_σ sufficiently large such that

$$\iint\limits_{(\text{supp }\varphi) \times (\mathbb{R}^N \setminus B_{R_\sigma})} |\mathcal{U}(x,y)|\, dxdy < \frac{\sigma}{2}.$$

Also, by definition of Ω_ε, we have $\Omega_\varepsilon \subset \text{supp }\varphi$ and $|\Omega_\varepsilon \times B_{R_\sigma}| \to 0$ as $\varepsilon \to 0^+$. Thus, since $\mathcal{U} \in L^1(\mathbb{R}^{2N})$, there exists $\delta_\sigma > 0$ and $\varepsilon_\sigma > 0$ such that for any $\varepsilon \in (0, \varepsilon_\sigma]$

$$|\Omega_\varepsilon \times B_{R_\sigma}| < \delta_\sigma \quad \text{and} \quad \iint\limits_{\Omega_\varepsilon \times B_{R_\sigma}} |\mathcal{U}(x,y)|\, dxdy < \frac{\sigma}{2}.$$

Therefore, for any $\varepsilon \in (0, \varepsilon_\sigma]$

$$\iint\limits_{\Omega_\varepsilon \times \mathbb{R}^N} |\mathcal{U}(x,y)|\, dxdy < \sigma,$$

from which we get

$$\lim_{\varepsilon \to 0^+} \iint\limits_{\Omega_\varepsilon \times \mathbb{R}^N} |\mathcal{U}(x,y)|\, dxdy = 0. \tag{3.17}$$

Combining (3.14) with (3.16), dividing by ε, letting $\varepsilon \to 0^+$ and using (3.17), we obtain

$$\|u_0\|^{2(\theta-1)} \langle u_0, \varphi \rangle - \lambda \int_\Omega (u_0^+)^{-\gamma} \varphi dx - \int_\Omega (u_0^+)^{2_s^*-1} \varphi dx \geq 0,$$

for any $\varphi \in X_0$. By the arbitrariness of φ, we prove that u_0 verifies (2.3), that is u_0 is a nontrivial solution of (2.2).

Finally, by considering $\varphi = u_0^-$ in (2.3) and using (3.12), we see that $\|u_0^-\| = 0$, which implies that u_0 is nonnegative. Moreover, by the maximum principle in [28, Proposition 2.2.8], we can deduce that u_0 is a positive solution of (2.2) and so also solves problem (1.1). This concludes the proof. □

We end this section observing that the result in Theorem 3.2 can be extended to more general Kirchhoff problems. That is, we can consider the following problem

$$\begin{cases} M\left(\iint\limits_{\mathbb{R}^{2N}} \dfrac{|u(x)-u(y)|^p}{|x-y|^{N+ps}} dxdy \right) (-\Delta)_p^s u = \dfrac{\lambda}{u^\gamma} + u^{p_s^*-1} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{in } \mathbb{R}^N \setminus \Omega, \end{cases} \tag{3.18}$$

where $p_s^* = pN/(N-ps)$, with here $N > ps$ and $p > 1$, while the Kirchhoff coefficient M satisfies condition

($\mathcal{M}$) $M : \mathbb{R}_0^+ \to \mathbb{R}_0^+$ *is continuous and nondecreasing. There exist numbers* $a > 0$ *and* ϑ *such that for any* $t \in \mathbb{R}_0^+$

$$\mathscr{M}(t) := \int_0^t M(\tau)d\tau \geq a\, t^{\vartheta}, \quad \text{with } \vartheta \begin{cases} \in (1, p_s^*/p) & \text{if } M(0) = 0, \\ = 1 & \text{if } M(0) > 0. \end{cases}$$

The main operator $(-\Delta)_p^s$ is the fractional p–Laplacian which may be defined, for any function $\varphi \in C_0^\infty(\Omega)$, as

$$(-\Delta)_p^s \varphi(x) = 2 \lim_{\varepsilon \to 0^+} \int_{\mathbb{R}^N \setminus B_\varepsilon(x)} \frac{|\varphi(x) - \varphi(y)|^{p-2}\big(\varphi(x) - \varphi(y)\big)}{|x-y|^{N+ps}} dy, \quad x \in \mathbb{R}^N,$$

where $B_\varepsilon(x) = \{y \in \mathbb{R}^N : \ |x-y| < \varepsilon\}$. Then, arguing as in the proof of Theorem 3.2 and observing that we have not used yet the assumption that $\partial\Omega$ is continuous, we can prove the following result.

Theorem 3.3. *Let* $s \in (0,1)$, $p > 1$, $N > ps$, $\gamma \in (0,1)$ *and let* Ω *be an open bounded subset of* $\mathbb{R}^N$. *Let* M *satisfy* ($\mathcal{M}$). *Then, there exists* $\lambda_0 > 0$ *such that for any* $\lambda \in (0, \lambda_0]$ *problem* (3.18) *admits a solution.*

4. A mountain pass solution for problem (2.4)

In this section we prove the existence of a positive solution for perturbed problem (2.4), by the mountain pass theorem. *For this, throughout this section we assume* $n \in \mathbb{N}$, *without further mentioning.* Now, we first prove that the related functional $J_{n,\lambda}$ satisfies all the geometric features required by the mountain pass theorem.

Lemma 4.1. *Let* $\rho \in (0,1]$, $\lambda_0 = \lambda_0(\rho) > 0$ *and* $\alpha = \alpha(\rho) > 0$ *be given as in Lemma 3.1. Then, for any* $\lambda \in (0, \lambda_0]$ *and any* $u \in X_0$, *with* $\|u\| \leq \rho$, *functional* $J_{n,\lambda}(u) \geq \alpha$.

Furthermore, there exists $e \in X_0$, *with* $\|e\| > \rho$, *such that* $J_{n,\lambda}(e) < 0$.

Proof. Since $\gamma \in (0,1)$, by the subadditivity of $t \mapsto t^{1-\gamma}$, we have

$$(u^+ + 1/n)^{1-\gamma} - (1/n)^{1-\gamma} \leq (u^+)^{1-\gamma} \quad \text{a.e. in } \Omega, \tag{4.1}$$

for any $u \in X_0$ and any $n \in \mathbb{N}$. Thus, we have $J_{n,\lambda}(u) \geq J_\lambda(u)$ for any $u \in X_0$ and the first part of lemma directly follows by Lemma 3.1.

For any $v \in X_0$, with $v^+ \not\equiv 0$, and $t > 0$ we have

$$J_{n,\lambda}(tv) = \frac{t^{2\theta}}{2\theta}\|v\|^{2\theta} - \frac{\lambda}{1-\gamma}\int_\Omega \left[(tv^+ + 1/n)^{1-\gamma} - (1/n)^{1-\gamma}\right] dx - \frac{t^{2_s^*}}{2_s^*}\|v^+\|_{2_s^*}^{2_s^*} \to -\infty \quad \text{as } t \to \infty,$$

being $1 - \gamma < 1 < 2\theta < 2_s^*$. Hence, we can find $e \in X_0$, with $\|e\| > \rho$ sufficiently large, such that $J_{n,\lambda}(e) < 0$. This concludes the proof. □

We discuss now the compactness property for the functional $J_{n,\lambda}$, given by the Palais–Smale condition. We recall that $\{u_k\}_k \subset X_0$ is a Palais–Smale sequence for $J_{n,\lambda}$ at level $c \in \mathbb{R}$ if

$$J_{n,\lambda}(u_k) \to c \quad \text{and} \quad J'_{n,\lambda}(u_k) \to 0 \quad \text{in } (X_0)' \text{ as } k \to \infty. \tag{4.2}$$

We say that $J_{n,\lambda}$ satisfies the Palais–Smale condition at level c if any Palais–Smale sequence $\{u_k\}_k$ at level c admits a convergent subsequence in X_0.

Before proving this compactness condition, we introduce the following positive constants which will help us for a better explanation

$$D_1 = \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right) S^{\frac{2_s^*\theta}{2_s^*-2\theta}} \qquad D_2 = \frac{\left[\left(\frac{1}{1-\gamma} + \frac{1}{2_s^*}\right) |\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}} S^{-\frac{1-\gamma}{2}}\right]^{\frac{2\theta}{2\theta-1+\gamma}}}{\left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)^{\frac{1-\gamma}{2\theta-1+\gamma}}}. \tag{4.3}$$

Lemma 4.2. *Let $\lambda > 0$. Then, the functional $J_{n,\lambda}$ satisfies the Palais–Smale condition at any level $c \in \mathbb{R}$ verifying*

$$c < D_1 - D_2 \lambda^{2\theta/(2\theta-1+\gamma)}, \tag{4.4}$$

with D_1, $D_2 > 0$ given as in (4.3).

Proof. Let $\lambda > 0$ and let $\{u_k\}_k$ be a Palais–Smale sequence in X_0 at level $c \in \mathbb{R}$, with c satisfying (4.4). We first prove the boundedness of $\{u_k\}_k$. By (4.2) there exists $\sigma > 0$ such that, as $k \to \infty$

$$\begin{aligned} c + \sigma\|u_k\| + o(1) &\geq J_{n,\lambda}(u_k) - \frac{1}{2_s^*}\langle J'_{n,\lambda}(u_k), u_k\rangle \\ &= \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\|u_k\|^{2\theta} - \frac{\lambda}{1-\gamma}\int_\Omega [(u_k^+ + 1/n)^{1-\gamma} - (1/n)^{1-\gamma}]dx \\ &\quad + \frac{\lambda}{2_s^*}\int_\Omega (u_k^+ + 1/n)^{-\gamma} u_k dx \\ &\geq \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\|u_k\|^{2\theta} - \lambda\left(\frac{1}{1-\gamma} + \frac{1}{2_s^*}\right)\int_\Omega |u_k|^{1-\gamma}dx \\ &\geq \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\|u_k\|^{2\theta} - \lambda\left(\frac{1}{1-\gamma} + \frac{1}{2_s^*}\right)|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}} S^{-\frac{1-\gamma}{2}}\|u_k\|^{1-\gamma} \end{aligned}$$

where the last two inequalities follow by (2.7), (4.1) and the Hölder inequality. Therefore, $\{u_k\}_k$ is bounded in X_0, being $1-\gamma < 1 < 2\theta$. Also, $\{u_k^-\}_k$ is bounded in X_0 and by (4.2) we have

$$\lim_{k\to\infty}\langle J'_{n,\lambda}(u_k), -u_k^-\rangle = \lim_{k\to\infty}\|u_k\|^{2(\theta-1)}\langle u_k, -u_k^-\rangle = 0.$$

Thus, by inequality (3.12) we deduce that $\|u_k^-\| \to 0$ as $k \to \infty$, which yields

$$J_{n,\lambda}(u_k) = J_{n,\lambda}(u_k^+) + o(1) \quad \text{and} \quad J'_{n,\lambda}(u_k) = J'_{n,\lambda}(u_k^+) + o(1) \quad \text{as } k \to \infty.$$

Hence, we can suppose that $\{u_k\}_k$ is a sequence of nonnegative functions.

By the boundedness of $\{u_k\}_k$ and by using [26, Lemma 8] and [6, Theorem 4.9], there exist a subsequence, still denoted by $\{u_k\}_k$, and a function $u \in X_0$ such that

$$\begin{aligned} &u_k \rightharpoonup u \text{ in } X_0, && \|u_k\| \to \mu, \\ &u_k \rightharpoonup u \text{ in } L^{2_s^*}(\Omega), && \|u_k - u\|_{2_s^*} \to \ell, \\ &u_k \to u \text{ in } L^p(\Omega) \text{ for any } p \in [1, 2_s^*), && u_k \to u \text{ a.e. in } \Omega, \quad u_k \leq h \text{ a.e. in } \Omega, \end{aligned} \tag{4.5}$$

as $k \to \infty$, with $h \in L^p(\Omega)$ for a fixed $p \in [1, 2_s^*)$. If $\mu = 0$, then immediately $u_k \to 0$ in X_0 as $k \to \infty$. Hence, let us assume that $\mu > 0$.

Since $n \in \mathbb{N}$, by (4.5) it follows that

$$\left|\frac{u_k - u}{(u_k + 1/n)^\gamma}\right| \leq n^\gamma(h + |u|) \quad \text{a.e. in } \Omega,$$

so by the dominated convergence theorem and (4.5), we have

$$\lim_{k\to\infty}\int_\Omega \frac{u_k-u}{(u_k+1/n)^\gamma}dx=0. \tag{4.6}$$

By (4.5) and [7, Theorem 2] we have

$$\|u_k\|^2=\|u_k-u\|^2+\|u\|^2+o(1),\quad \|u_k\|_{2_s^*}^{2_s^*}=\|u_k-u\|_{2_s^*}^{2_s^*}+\|u\|_{2_s^*}^{2_s^*}+o(1) \tag{4.7}$$

as $k\to\infty$. Consequently, we deduce from (4.2), (4.5), (4.6) and (4.7) that, as $k\to\infty$

$$\begin{aligned} o(1)&=\langle J'_{n,\lambda}(u_k),u_k-u\rangle\\ &=\|u_k\|^{2(\theta-1)}\langle u_k,u_k-u\rangle-\lambda\int_\Omega\frac{u_k-u}{(u_k+1/n)^\gamma}dx-\int_\Omega u_k^{2_s^*-1}(u_k-u)dx\\ &=\mu^{2(\theta-1)}(\mu^2-\|u_k\|^2)-\|u_k\|_{2_s^*}^{2_s^*}+\|u\|_{2_s^*}^{2_s^*}+o(1)=\mu^{2(\theta-1)}\|u_k-u\|^2-\|u_k-u\|_{2_s^*}^{2_s^*}+o(1). \end{aligned}$$

Therefore, we have proved the crucial formula

$$\mu^{2(\theta-1)}\lim_{k\to\infty}\|u_k-u\|^2=\lim_{k\to\infty}\|u_k-u\|_{2_s^*}^{2_s^*}. \tag{4.8}$$

If $\ell=0$, since $\mu>0$, by (4.5) and (4.8) we have $u_k\to u$ in X_0 as $k\to\infty$, concluding the proof.

Thus, let us assume by contradiction that $\ell>0$. By (2.7), the notation in (4.5) and (4.8), we get

$$\ell^{2_s^*}\geq S\,\mu^{2(\theta-1)}\ell^2. \tag{4.9}$$

Noting that (4.8) implies in particular that

$$\mu^{2(\theta-1)}\big(\mu^2-\|u\|^2\big)=\ell^{2_s^*},$$

using (4.9), it follows that

$$\big(\ell^{2_s^*}\big)^{2s/N}=(\mu^{2(\theta-1)})^{2s/N}\big(\mu^2-\|u\|^2\big)^{2s/N}\geq S\,\mu^{2(\theta-1)}.$$

From this, we obtain

$$\mu^{4s/N}\geq\big(\mu^2-\|u\|^2\big)^{2s/N}\geq S\,(\mu^{2(\theta-1)})^{\frac{N-2s}{N}},$$

and considering $N<2s\theta/(\theta-1)=2s\theta'$, we have

$$\mu^2\geq S^{\frac{N}{2s\theta-N(\theta-1)}}. \tag{4.10}$$

Indeed, the restriction $N/(2\theta')<s$ follows directly from the fact that $1<\theta<2_s^*/2=N/(N-2s)$. By (4.1) and considering that $n\in\mathbb{N}$, for any $k\in\mathbb{N}$ we have

$$J_{n,\lambda}(u_k)-\frac{1}{2_s^*}\left\langle J'_{n,\lambda}(u_k),u_k\right\rangle\geq\left(\frac{1}{2\theta}-\frac{1}{2_s^*}\right)\|u_k\|^{2\theta}-\lambda\left(\frac{1}{1-\gamma}+\frac{1}{2_s^*}\right)\int_\Omega u_k^{1-\gamma}dx.$$

From this, as $k \to \infty$, being $\theta \geq 1$, by (4.2), (4.5), (4.7), (4.10), the Hölder inequality and the Young inequality, we obtain

$$\begin{aligned} c &\geq \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\left(\mu^{2\theta} + \|u\|^{2\theta}\right) - \lambda\left(\frac{1}{1-\gamma} + \frac{1}{2_s^*}\right)|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}} S^{-\frac{1-\gamma}{2}} \|u\|^{1-\gamma} \\ &\geq \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\left(\mu^{2\theta} + \|u\|^{2\theta}\right) - \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\|u\|^{2\theta} \\ &\quad - \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)^{-\frac{1-\gamma}{2\theta-1+\gamma}} \left[\lambda\left(\frac{1}{1-\gamma} + \frac{1}{2_s^*}\right)|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}} S^{-\frac{1-\gamma}{2}}\right]^{\frac{2\theta}{2\theta-1+\gamma}} \\ &\geq \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right) S^{\frac{2_s^*\theta}{2_s^*-2\theta}} - \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)^{-\frac{1-\gamma}{2\theta-1+\gamma}} \left[\lambda\left(\frac{1}{1-\gamma} + \frac{1}{2_s^*}\right)|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}} S^{-\frac{1-\gamma}{2}}\right]^{\frac{2\theta}{2\theta-1+\gamma}} \end{aligned}$$

which contradicts (4.4), since (4.3). This concludes the proof. □

We now give a control from above for functional $J_{n,\lambda}$, under a suitable situation. For this, we assume, without loss of generality, that $0 \in \Omega$. By [10] we know that the infimum in (2.7) is attained at the function

$$u_\varepsilon(x) = \frac{\varepsilon^{(N-2s)/2}}{(\varepsilon^2 + |x|^2)^{(N-2s)/2}}, \quad \text{with } \varepsilon > 0, \tag{4.11}$$

that is, it holds true that

$$\iint_{\mathbb{R}^{2N}} \frac{|u_\varepsilon(x) - u_\varepsilon(y)|^2}{|x-y|^{N+2s}} dxdy = S\|u_\varepsilon\|^2_{L^{2_s^*}(\mathbb{R}^N)}.$$

Let us fix $r > 0$ such that $B_{4r} \subset \Omega$, where $B_{4r} = \{x \in \mathbb{R}^N : |x| < 4r\}$, and let us introduce a cut–off function $\phi \in C^\infty(\mathbb{R}^N, [0,1])$ such that

$$\phi = \begin{cases} 1 & \text{in } B_r, \\ 0 & \text{in } \mathbb{R}^N \setminus B_{2r}. \end{cases} \tag{4.12}$$

For any $\varepsilon > 0$, we set

$$\psi_\varepsilon = \frac{\phi u_\varepsilon}{\|\phi u_\varepsilon\|^2_{2_s^*}} \in X_0. \tag{4.13}$$

Then, we can prove the following result.

Lemma 4.3. *There exist $\psi \in X_0$ and $\lambda_1 > 0$ such that for any $\lambda \in (0, \lambda_1)$*

$$\sup_{t \geq 0} J_{n,\lambda}(t\psi) < D_1 - D_2 \lambda^{2\theta/(2\theta-1+\gamma)},$$

with D_1, $D_2 > 0$ given as in (4.3).

Proof. Let λ, $\varepsilon > 0$. Let u_ε and ψ_ε be as respectively in (4.11) and in (4.13). By (2.6), we have $J_{n,\lambda}(t\psi_\varepsilon) \to -\infty$ as $t \to \infty$, so that there exists $t_\varepsilon > 0$ such that $J_{n,\lambda}(t_\varepsilon \psi_\varepsilon) = \max_{t \geq 0} J_{n,\lambda}(t\psi_\varepsilon)$. By Lemma 4.1 we know that $J_{n,\lambda}(t_\varepsilon \psi_\varepsilon) \geq \alpha > 0$. Hence, by continuity of $J_{n,\lambda}$, there exist two numbers t_0, $t^* > 0$ such that $t_0 \leq t_\varepsilon \leq t^*$.

Now, since $\|u_\varepsilon\|_{L^{2_s^*}(\mathbb{R}^N)}$ is independent of ε, by [27, Proposition 21] we have

$$\|\psi_\varepsilon\|^2 \le \frac{\iint_{\mathbb{R}^{2N}} \frac{|u_\varepsilon(x)-u_\varepsilon(y)|^2}{|x-y|^{N+2s}}dxdy}{\|\phi u_\varepsilon\|_{2_s^*}^2} = S + O(\varepsilon^{N-2s}),$$

from which, by the elementary inequality

$$(a+b)^p \le a^p + p(a+1)^{p-1}b, \quad \text{for any } a>0,\ b\in[0,1],\ p\ge 1,$$

with $p=2\theta$, it follows that, as $\varepsilon \to 0^+$

$$\|\psi_\varepsilon\|^{2\theta} \le S^\theta + O(\varepsilon^{N-2s}).$$

Hence, by the last inequality, (2.6) and being $t_0 \le t_\varepsilon \le t^*$, for any $\varepsilon>0$ sufficiently small, we have

$$J_{n,\lambda}(t_\varepsilon\psi_\varepsilon) \le \frac{t^{2\theta}}{2\theta}S^\theta + C_1\varepsilon^{N-2s} - \frac{\lambda}{1-\gamma}\int_\Omega [(t_0\psi_\varepsilon + 1/n)^{1-\gamma} - (1/n)^{1-\gamma}]dx - \frac{t^{2_s^*}}{2_s^*}, \tag{4.14}$$

with C_1 a suitable positive constant. We observe that

$$\max_{t\ge 0}\left(\frac{t^{2\theta}}{2\theta}S^\theta - \frac{t^{2_s^*}}{2_s^*}\right) = \left(\frac{1}{2\theta}-\frac{1}{2_s^*}\right)S^{\frac{2_s^*\theta}{2_s^*-2\theta}}.$$

Thus, by (4.14) it follows that

$$J_{n,\lambda}(t_\varepsilon\psi_\varepsilon) \le \left(\frac{1}{2\theta}-\frac{1}{2_s^*}\right)S^{\frac{2_s^*\theta}{2_s^*-2\theta}} + C_1\varepsilon^{N-2s} - \frac{\lambda}{1-\gamma}\int_\Omega [(t_0\psi_\varepsilon + 1/n)^{1-\gamma} - (1/n)^{1-\gamma}]dx. \tag{4.15}$$

Now, let us consider a positive number q satisfying

$$\frac{(N-2s)(1-\gamma)-2q(N-2s)(1-\gamma)+2_s^*qN}{2_s^*}\cdot\frac{2\theta}{2\theta-1+\gamma}\cdot\frac{1}{N-2s}+1-\frac{2\theta}{2\theta-1+\gamma}<0, \tag{4.16}$$

that is, being $2<2\theta<2_s^*$, $N>2s$ and $\gamma\in(0,1)$, such that

$$0<q<\frac{(N-2s)(2_s^*-2\theta)(1-\gamma)}{2\theta N(2_s^*-2)+4\theta N\gamma+8\theta s(1-\gamma)}.$$

By the elementary inequality

$$a^{1-\gamma}-(a+b)^{1-\gamma} \le -(1-\gamma)b^{\frac{1-\gamma}{p}}a^{\frac{(p-1)(1-\gamma)}{p}}, \quad \text{for any } a>0,\ b>0 \text{ large enough},\ p>1,$$

with $p=2_s^*/2$, and considering $\varepsilon<r^{1/q}$ sufficiently small, with r given by (4.12), we have

$$\begin{aligned}
&-\frac{1}{1-\gamma}\int_{\{x\in\Omega:\,|x|\le\varepsilon^q\}}[(t_0\psi_\varepsilon+1/n)^{1-\gamma}-(1/n)^{1-\gamma}]dx\\
&\le -\widetilde{C}\varepsilon^{\frac{(N-2s)(1-\gamma)}{2_s^*}}\int_{\{x\in\Omega:\,|x|\le\varepsilon^q\}}\left[\frac{1}{(|x|^2+\varepsilon^2)^{\frac{N-2s}{2}}}\right]^{\frac{2(1-\gamma)}{2_s^*}}dx\\
&\le -C_2\varepsilon^{\frac{(N-2s)(1-\gamma)-2q(N-2s)(1-\gamma)+2_s^*qN}{2_s^*}},
\end{aligned} \tag{4.17}$$

with $\widetilde{C}$ and C_2 two positive constant independent by ε. By combining (4.15) with (4.17), we get

$$J_{n,\lambda}(t_\varepsilon\psi_\varepsilon) \le \left(\frac{1}{2\theta}-\frac{1}{2_s^*}\right)S^{\frac{2_s^*\theta}{2_s^*-2\theta}} + C_1\varepsilon^{N-2s} - C_2\lambda\varepsilon^{\frac{(N-2s)(1-\gamma)-2q(N-2s)(1-\gamma)+2_s^*qN}{2_s^*}}. \tag{4.18}$$

Thus, let us consider $\lambda^* > 0$ such that

$$D_1 - D_2\lambda^{\frac{2\theta}{2\theta-1+\gamma}} > 0 \quad \text{for any } \lambda \in (0, \lambda^*)$$

and let us set

$$\nu_1 = \frac{2q\theta}{(2\theta-1+\gamma)(N-2s)}, \quad \nu_2 = \frac{2\theta[(N-2s)(1-\gamma) - 2q(N-2s)(1-\gamma) + 2_s^* qN]}{2_s^*(2\theta-1+\gamma)(N-2s)} + 1,$$
$$\nu_3 = \nu_2 - \frac{2\theta}{2\theta-1+\gamma}, \quad \lambda_1 = \min\left\{\lambda^*, r^{1/\nu_1}, \left(\frac{C_2}{C_1+D_2}\right)^{-1/\nu_3}\right\},$$

where r and q are given respectively in (4.12) and (4.16), while we still consider D_1 and D_2 as defined in (4.3). Then, by considering $\varepsilon = \lambda^{\nu_1/q}$ in (4.18), since (4.16) implies that $\nu_3 < 0$, for any $\lambda \in (0, \lambda_1)$ we have

$$J_{n,\lambda}(t_\varepsilon\psi_\varepsilon) \le D_1 + C_1\lambda^{\frac{2\theta}{2\theta-1+\gamma}} - C_2\lambda^{\nu_2} = D_1 + \lambda^{\frac{2\theta}{2\theta-1+\gamma}}(C_1 - C_2\lambda^{\nu_3}) < D_1 - D_2\lambda^{\frac{2\theta}{2\theta-1+\gamma}},$$

which concludes the proof. □

We can now prove the existence result for (2.4) by applying the mountain pass theorem.

Theorem 4.4. *There exists $\overline{\lambda} > 0$ such that, for any $\lambda \in (0, \overline{\lambda})$, problem (2.4) has a positive solution $v_n \in X_0$, with*

$$\alpha < J_{n,\lambda}(v_n) < D_1 - D_2\lambda^{2\theta/(2\theta-1+\gamma)}, \tag{4.19}$$

where α, D_1 and D_2 are given respectively in Lemma 3.1 and (4.3).

Proof. Let $\overline{\lambda} = \min\{\lambda_0, \lambda_1\}$, with λ_0 and λ_1 given respectively in Lemmas 3.1 and 4.3. Let us consider $\lambda \in (0, \overline{\lambda})$. By Lemma 4.1, functional $J_{n,\lambda}$ verifies the mountain pass geometry. For this, we can set the critical mountain pass level as

$$c_{n,\lambda} = \inf_{g\in\Gamma}\max_{t\in[0,1]} J_{n,\lambda}(g(t)), \tag{4.20}$$

where

$$\Gamma = \{g \in C([0,1], X_0) : \ g(0) = 0, \ J_{n,\lambda}(g(1)) < 0\}.$$

By Lemmas 4.1 and 4.3, we get

$$0 < \alpha < c_{n,\lambda} \le \sup_{t\ge 0} J_{n,\lambda}(t\psi) < D_1 - D_2\lambda^{2\theta/(2\theta-1+\gamma)}.$$

Hence, by Lemma 4.2 functional $J_{n,\lambda}$ satisfies the Palais–Smale condition at level $c_{n,\lambda}$. Thus, the mountain pass theorem gives the existence of a critical point $v_n \in X_0$ for $J_{n,\lambda}$ at level $c_{n,\lambda}$. Since $J_{n,\lambda}(v_n) = c_{n,\lambda} > \alpha > 0 = J_{n,\lambda}(0)$, v_n is a nontrivial solution of (2.4). Furthermore, by (2.5) with test function $\varphi = v_n^-$ and inequality (3.12), we can see that $\|v_n^-\| = 0$, which implies v_n is nonnegative. By the maximum principle in [28, Proposition 2.2.8], we have that v_n is a positive solution of (2.4), concluding the proof. □

5. A second solution for problem (1.1)

In this last section we prove the existence of a second solution for problem (1.1), as a limit of solutions of the perturbed problem (2.4). For this, here we need the assumption that $\partial\Omega$ is continuous, in order to apply a density argument for space X_0.

Proof of Theorem 1.1. Let us consider $\overline{\lambda}$ as given in Theorem 4.4 and let $\lambda \in (0, \overline{\lambda})$. Since $\overline{\lambda} \leq \lambda_0$, by Theorem 3.2 we know that problem (1.1) admits a solution u_0 with $J_\lambda(u_0) < 0$.

In order to find a second solution for (1.1), let $\{v_n\}_n$ be a family of positive solutions of (2.4). By (2.7), (4.1), (4.19) and the Hölder inequality, we have

$$\begin{aligned} D_1 - D_2\lambda^{\frac{2\theta}{2\theta-1+\gamma}} &> J_{n,\lambda}(v_n) - \frac{1}{2_s^*}\langle J'_{n,\lambda}(v_n), v_n\rangle \\ &= \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\|v_n\|^{2\theta} - \frac{\lambda}{1-\gamma}\int_\Omega [(v_n + 1/n)^{1-\gamma} - (1/n)^{1-\gamma}]dx \\ &\quad + \frac{\lambda}{2_s^*}\int_\Omega (v_n + 1/n)^{-\gamma} v_n dx \\ &\geq \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\|v_n\|^{2\theta} - \frac{\lambda}{1-\gamma}\int_\Omega v_n^{1-\gamma} dx \\ &\geq \left(\frac{1}{2\theta} - \frac{1}{2_s^*}\right)\|v_n\|^{2\theta} - \frac{\lambda}{1-\gamma}|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}} S^{-\frac{1-\gamma}{2}}\|v_n\|^{1-\gamma}, \end{aligned}$$

which yields that $\{v_n\}_n$ is bounded in X_0, being $1-\gamma < 1 < 2\theta$. Hence, by using [26, Lemma 8] and [6, Theorem 4.9], there exist a subsequence, still denoted by $\{v_n\}_n$, and a function $v_0 \in X_0$ such that

$$\begin{array}{ll} v_n \rightharpoonup v_0 \text{ in } X_0, & \|v_n\| \to \mu, \\ v_n \rightharpoonup v_0 \text{ in } L^{2_s^*}(\Omega), & \|v_n - v_0\|_{2_s^*} \to \ell, \\ v_n \to v_0 \text{ in } L^p(\Omega) \text{ for any } p \in [1, 2_s^*), & v_n \to v_0 \text{ a.e. in } \Omega. \end{array} \tag{5.1}$$

We want to prove that $v_n \to v_0$ in X_0 as $n \to \infty$. When $\mu = 0$, by (5.1) we have $v_n \to 0$ in X_0 as $n \to \infty$. For this, we suppose $\mu > 0$. We observe that

$$0 \leq \frac{v_n}{(v_n + 1/n)^\gamma} \leq v_n^{1-\gamma} \quad \text{a.e. in } \Omega,$$

so by the Vitali convergence theorem and (5.1), it follows that

$$\lim_{n\to\infty}\int_\Omega \frac{v_n}{(v_n + 1/n)^\gamma} dx = \int_\Omega v_0^{1-\gamma} dx. \tag{5.2}$$

By using (2.5) for v_n and test function $\varphi = v_n$, by (5.1) and (5.2), as $n \to \infty$ we have

$$\mu^{2\theta} - \lambda\int_\Omega v_0^{1-\gamma} dx + \|v_n\|_{2_s^*}^{2_s^*} = o(1). \tag{5.3}$$

For any $n \in \mathbb{N}$, by an immediate calculation in (2.4) we see that

$$\|v_n\|^{2\theta}(-\Delta)^s v_n \geq \min\left\{1, \frac{\lambda}{2^\gamma}\right\} \quad \text{in } \Omega.$$

Thus, since $\{v_n\}_n$ is bounded in X_0 and by using a standard comparison argument (see [4, Lemma 2.1]) and the maximum principle in [28, Proposition 2.2.8], for any $\widetilde{\Omega} \subset\subset \Omega$, there exists a constant $c_{\widetilde{\Omega}} > 0$ such that

$$v_n \geq c_{\widetilde{\Omega}} > 0, \quad \text{a.e. in } \widetilde{\Omega} \text{ and for any } n \in \mathbb{N}. \tag{5.4}$$

Now, let $\varphi \in C_0^\infty(\Omega)$, with supp $\varphi = \widetilde{\Omega} \subset\subset \Omega$. By (5.4) we have

$$0 \leq \left|\frac{\varphi}{(v_n + 1/n)^\gamma}\right| \leq \frac{|\varphi|}{c_{\widetilde{\Omega}}^\gamma} \quad \text{a.e. in } \Omega,$$

so that by (5.1) and the dominated convergence theorem

$$\lim_{n\to\infty}\int_\Omega \frac{\varphi}{(v_n+1/n)^\gamma}dx=\int_\Omega v_0^{-\gamma}\varphi dx. \tag{5.5}$$

Thus, by considering (2.5) for v_n, sending $n\to\infty$ and using (5.1) and (5.5), for any $\varphi\in C_0^\infty(\Omega)$ it follows that

$$\mu^{2(\theta-1)}\langle v_0,\varphi\rangle-\lambda\int_\Omega v_0^{-\gamma}\varphi dx+\int_\Omega v_0^{2_s^*-1}\varphi dx=0. \tag{5.6}$$

However, since $\partial\Omega$ is continuous, by [15, Theorem 6] the space $C_0^\infty(\Omega)$ is dense in X_0. Thus, by a strandard density argument, (5.6) holds true for any $\varphi\in X_0$. By combining (5.3) and (5.6) with test function $\varphi=v_0$, as $n\to\infty$ we get

$$\mu^{2(\theta-1)}(\mu^2-\|v_0\|^2)=\|v_n\|_{2_s^*}^{2_s^*}-\|v_0\|_{2_s^*}^{2_s^*}+o(1)$$

and by (5.1) and [7, Theorem 2] we have

$$\mu^{2(\theta-1)}\lim_{n\to\infty}\|v_n-v_0\|^2=\ell^{2_s^*}. \tag{5.7}$$

If $\ell=0$, then $v_n\to v_0$ in X_0 as $n\to\infty$, since $\mu>0$.

Let us suppose $\ell>0$ by contradiction. Arguing as in Lemma 4.2, by (5.1) and (5.7) we get (4.10). Therefore, being $\theta\geq 1$, by (4.1), (4.10), (4.19), (5.1), the Hölder inequality and the Young inequality, we have

$$\begin{aligned}
D_1-D_2\lambda^{\frac{2\theta}{2\theta-1+\gamma}} &> J_{n,\lambda}(v_n)-\frac{1}{2_s^*}\left\langle J'_{n,\lambda}(v_n),v_n\right\rangle\\
&\geq\left(\frac{1}{2\theta}-\frac{1}{2_s^*}\right)\left(\mu^{2\theta}+\|v_0\|^{2\theta}\right)-\lambda\left(\frac{1}{1-\gamma}+\frac{1}{2_s^*}\right)|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}}S^{-\frac{1-\gamma}{2}}\|v_0\|^{1-\gamma}\\
&\geq\left(\frac{1}{2\theta}-\frac{1}{2_s^*}\right)S^{\frac{2_s^*\theta}{2_s^*-2\theta}}\\
&\quad-\left(\frac{1}{2\theta}-\frac{1}{2_s^*}\right)^{-\frac{1-\gamma}{2\theta-1+\gamma}}\left[\lambda\left(\frac{1}{1-\gamma}+\frac{1}{2_s^*}\right)|\Omega|^{\frac{2_s^*-1+\gamma}{2_s^*}}S^{-\frac{1-\gamma}{2}}\right]^{\frac{2\theta}{2\theta-1+\gamma}}
\end{aligned}$$

which is the desired contradiction, thanks to (4.3).

Therefore, $v_n\to v_0$ in X_0 as $n\to\infty$ and by (2.3) and (2.5) we immediately see that v_0 is a solution of problem (2.2). Furthermore, by (4.19) we have $J_\lambda(v_0)\geq\alpha>0$, which also implies that v_0 is nontrivial. Reasoning as at the end of the proof of Theorem 4.4, we conclude that v_0 is a positive solution of (2.2) and so v_0 also solves problem (1.1). Finally, v_0 is different from u_0, since $J_\lambda(v_0)>0>J_\lambda(u_0)$. □

Acknowledgments

The author is supported by *Coordenação de Aperfeiçonamento de pessoal de nível superior* through the fellowship PNPD–CAPES 33003017003P5. The author is member of the *Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni* (GNAMPA) of the *Istituto Nazionale di Alta Matematica "G. Severi"* (INdAM).

References

[1] B. Abdellaoui, M. Medina, I. Peral and A. Primo, *The effect of the Hardy potential in some Calderón–Zygmund properties for the fractional Laplacian*, J. Differential Equations 260, 8160–8206 (2016).

[2] G. Autuori, A. Fiscella and P. Pucci, *Stationary Kirchhoff problems involving a fractional elliptic operator and a critical nonlinearity*, Nonlinear Anal. 125, 699–714 (2015).

[3] B. Barrios, I. De Bonis, M. Medina and I. Peral, *Semilinear problems for the fractional laplacian with a singular nonlinearity*, Open Math. 13, 390–407 (2015).

[4] B. Barrios, M. Medina and I. Peral, *Some remarks on the solvability of non local elliptic problems with the Hardy potential*, Contemp. Math. 16, 29 pp. (2013).

[5] Z. Binlin, A. Fiscella and S. Liang, *Infinitely many solutions for critical degenerate Kirchhoff type equations involving the fractional p–Laplacian*, submitted for publication.

[6] H. Brézis, "Functional analysis, Sobolev spaces and partial differential equations", *Universitext*, Springer, New York, 2011, xiv+599 pp.

[7] H. Brézis and E. Lieb, *A relation between pointwise convergence of functions and convergence of functionals*, Proc. Amer. Math. Soc. 88, 486–490 (1983).

[8] A. Canino, L. Montoro, B. Sciunzi and M. Squassina, *Nonlocal problems with singular nonlinearity*, to appear on Bull. Sci. Math. (2017).

[9] M. Caponi and P. Pucci, *Existence theorems for entire solutions of stationary Kirchhoff fractional p–Laplacian equations*, Ann. Mat. Pura Appl. 195, 2099–2129 (2016).

[10] A. Cotsiolis and N. Tavoularis, *Best constants for Sobolev inequalities for higher order fractional derivatives*, J. Math. Anal. Appl. 295, 225–236 (2004).

[11] E. Di Nezza, G. Palatucci and E. Valdinoci, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math. 136, 521–573 (2012).

[12] A. Fiscella, *Infinitely many solutions for a critical Kirchhoff type problem involving a fractional operator*, Differ. Integral Equations 29, 513–530 (2016).

[13] A. Fiscella and P. Pucci, *Kirchhoff Hardy fractional problems with lack of compactness*, submitted for pubblication.

[14] A. Fiscella and P. Pucci, *p–fractional Kirchhoff equations involving critical nonlinearities*, Nonlinear Anal. Real World Appl. 35, 350–378 (2017).

[15] A. Fiscella, R. Servadei and E. Valdinoci, *Density properties for fractional Sobolev spaces*, Ann. Acad. Sci. Fenn. Math. 40, 235–253 (2015).

[16] A. Fiscella and E. Valdinoci, *A critical Kirchhoff type problem involving a nonlocal operator*, Nonlinear Anal. 94, 156–170 (2014).

[17] C.–Y. Lei, J.–F. Liao and C.–L. Tang, *Multiple positive solutions for Kirchhoff type of problems with singularity and critical exponents*, J. Math. Anal. Appl. 421, 521–538 (2015).

[18] J.–F. Liao, X.–F. Ke, C.–Y. Lei and C.–L. Tang, *A uniqueness result for Kirchhoff type problems with singularity*, Appl. Math. Lett. 59, 24–30 (2016).

[19] J.–F. Liao, P. Zhang, J. Liu and C.–L. Tang, *Existence and multiplicity of positive solutions for a class of Kirchhoff type problems with singularity*, J. Math. Anal. Appl. 430, 1124–1148 (2015).

[20] X. Liu and Y. Sun, *Multiple positive solutions for Kirchhoff type of problems with singularity*, Commun. Pure Appl. Anal. 12, 721–733 (2013).

[21] R.–Q. Liu, C.–L. Tang, J.–F. Liao and X.–P. Wu, *Positive solutions of Kirchhoff type problem with singular and critical nonlinearities in dimension four*, Commun. Pure Appl. Anal. 15, 1841–1856 (2016).

[22] G. Molica Bisci, V. Rădulescu and R. Servadei, "Variational Methods for Nonlocal Fractional Problems", Encyclopedia of Mathematics and its Applications, 162, Cambride University Press, Cambridge, 2016, xvi+383 pp.

[23] P. Piersanti and P. Pucci, *Entire solutions for critical p–fractional Hardy Schrödinger Kirchhoff equations*, to appear on Publ. Mat. (2017).

[24] P. Pucci and S. Saldi, *Critical stationary Kirchhoff equations in $\mathbb{R}^N$ involving nonlocal operators*, Rev. Mat. Iberoam. 32, 1–22 (2016).

[25] P. Pucci, M. Xiang and B. Zhang, *Existence and multiplicity of entire solutions for fractional p–Kirchhoff equations*, Adv. Nonlinear Anal. 5, 27–55 (2016).

[26] R. Servadei and E. Valdinoci, *Mountain Pass solutions for non-local elliptic operators*, J. Math. Anal. Appl. 389, 887–898 (2012).

[27] R. Servadei and E. Valdinoci, *The Brezis-Nirenberg result for the fractional Laplacian*, Trans. Amer. Math. Soc. 367, 67–102 (2015).
[28] L. Silvestre, *Regularity of the obstacle problem for a fractional power of the Laplace operator*, Comm. Pure Appl. Math. 60, 67–112 (2007).
[29] M. Xiang, B. Zhang and H. Qiu, *Existence of solutions for a critical fractional Kirchhoff type problem in* $\mathbb{R}^N$, to appear on Sci. China Math. (2017).

Alessio Fiscella, Departamento de Matemática, Universidade Estadual de Campinas, IMECC, Rua Sérgio Buarque de Holanda, 651, Campinas, SP CEP 13083–859 Brazil
E-mail address: fiscella@ime.unicamp.br